

Atomic Design of Three-Dimensional Photonic Z_2 Dirac and Weyl Points

HaiXiao Wang,¹ Lin Xu,¹ HuanYang Chen,^{1,*} and Jian-Hua Jiang^{1,†}

¹*College of Physics, Optoelectronics and Energy,*

& Collaborative Innovation Center of Suzhou Nano Science and Technology,

Soochow University, 1 Shizi Street, Suzhou 215006, China

(Dated: January 27, 2016)

arXiv:1601.02276v2 [cond-mat.mtrl-sci] 26 Jan 2016

*Electronic address: chy@suda.edu.cn

†Electronic address: joejhjiang@sina.com

Topological nodal points such as Dirac and Weyl points in photonic spectrum, as monopoles of the synthetic gauge fields in $\vec{k}$ -space, offer unique abilities of manipulating light. However, designing topological nodal points in photonic crystals is much more difficult than in electronic band materials due to lack of the atomic picture. Here we propose an atomic approach for the design of three-dimensional Dirac and Weyl points via Mie resonances which can be regarded as photonic local orbits, using hollow-cylinder hexagonal photonic crystal as an example. We discover a new type of topological degeneracy, the Z_2 Dirac points, which are monopoles of the $SU(2)$ Berry-flux. Our study provides effective methodology as well as a new prototype of topological nodal points for future topological photonics and applications.

Stimulated by the discovery of topological insulators[1, 2] and topological semimetals[3, 4], the study of topological nodal points in electronic energy spectrum has attracted a lot of attention in the past decade. Recently, research interest in topological phenomena is extended to atomic[5, 6], photonic[7–13], and acoustic[14–16] systems. Pseudo-spin $S = 1$ Dirac cone was found in two-dimensional (2D) photonic crystals[17], offering unique properties for manipulation of light via, e.g., zero refractive index[17], Klein tunneling[18], and enhanced single-mode lasing[19]. Lately, Weyl points are predicted[8, 12] and observed[11, 12] in three-dimensional (3D) photonic crystals. It is noticed that 3D Weyl points can provide effective angle- and frequency-selective transmission[20]. These developments, alongside with the progresses in 2D[21–25] and 3D[26] photonic topological insulators, are unfolding a revolutionary platform of topological photonics where propagation of light can be controlled via topology-induced surface states and synthetic Berry gauge fields for photons to realize unprecedented applications.

The manifestation of topology and Berry phases in electronic and photonic systems usually comes along with symmetry protection. Electronic topological insulators, for instance, have nontrivial topology protected by the time-reversal symmetry. In electronic and other fermionic systems, this protection is because $\mathcal{T}^2 = -1$ where $\mathcal{T}$ is the time-reversal operator. Thus time-reversal operation enables and protects double degeneracy and the appearance of helical edge states. For photons and other bosons, $\mathcal{T}^2 = 1$ forbids emergence of topological insulators due to lack of such double degeneracy. Thus in photonic systems topological insulators emerge only in systems with additional symmetries, such as rotation symmetry[25]

and nonsymmorphic symmetry[26].

However, due to the complexity of photonic energy bands, up till now the design of topological properties in photonic crystals is quite difficult particularly for three-dimensional photonic crystals. This is due to the essential difference between electronic and photonic energy bands: The photonic energy bands are consequences of multiple Bragg scattering of the vectorial electromagnetic waves as no dielectric material can trap light[27–29]. In contrast, electronic band structure can mostly be understood as hybridization of local atomic orbits. For materials with inversion symmetry, the Z_2 topological index can be calculated simply by counting band (parity) inversion at high symmetry points in the Brillouin zone. Such a simplified picture is not available in photonic crystals, creating lots of obstacles in designing and understanding the topological properties of photonic bands. Nevertheless, it was noted that Mie resonances play similar roles in understanding photonic energy bands as that of atomic orbits for electronic band structures[30, 31].

In this work we propose a simple design to realize 3D Dirac and Weyl points in photonic crystals made of dielectrics with isotropic permittivity. Our design is based on that the Mie resonances of the (isolated) hollow cylinders can be regarded as photonic local orbits. The photonic energy bands can be understood as derived from photon hopping between Mie resonances in neighboring unit cells[30, 31] (except for the lowest two photonic bands which also consist of the plane-wave component, see Supplementary Information). The hollow cylinder supports p - and d -wave Mie resonances with double degeneracy as protected by the C_{6v} and inversion symmetries[24, 25]. Exploiting such double degeneracy a pseudo-time-reversal operation $\mathcal{T}_p = i\hat{\sigma}_y\mathcal{T}$ with $\mathcal{T}_p^2 = -1$ can be defined where $\hat{\sigma}_y$ is the pseudo-spin Pauli matrix[25] (the double degenerate Hilbert space is defined as the pseudo-spin). This degeneracy property has been exploited to construct 2D photonic topological insulators recently[25, 32]. In our closely-packed hollow cylinder photonic crystal the p - and d -like bands can be viewed as derived from the p - and d -wave Mie resonances. A pair of unavoidable, accidental degeneracy points of the p - and d - photonic bands are discovered, which are identified as the topological Dirac points protected by the parity-time symmetry[33]. The Dirac points found here are the Z_2 monopoles of the $SU(2)$ Berry-flux. They are quite different from the Weyl points in photonic crystals[8, 12] which are monopoles of the $U(1)$ Berry-flux and have a Z topological charge. Our work is the first proposal of the Z_2 topological Dirac points in photonic crystals.

The hexagonal photonic crystal consists of hollow cylinders (with outer radius R_{out} and inner radius R_{in}) connected by micropillars (see Fig. 1). The height of each unit cell is $h = 0.6a$ with a being the lattice constant in the x - y plane. The micropillars are of the same height $0.2a$ and diameter $0.1a$. There are six micropillars in each unit cell (their positions are illustrated in Fig. 1b). The height of each hollow cylinder is $0.4a$. The image of the bulk photonic crystal is depicted in Fig. 1c. The Brillouin zone and high symmetry points are shown in Fig. 1d.

Long hollow cylinders support local electromagnetic (Mie) resonances of s , p , d , f ... symmetries (see Supplementary Information). In hexagonal lattice with C_{6v} symmetry, only the $L_z = 0, 1, 2, 3$ (i.e., s , p , d , and f) orbits are distinguishable, higher angular momentum orbits are mixed with those lower ones. Along the z direction, the photonic “wavefunction” can have zero, one, or multiple nodes. The wavefunctions of the photonic bands of interest (the p - and d - bands) have zero nodes, i.e., they are *parity-even* along the z direction. The micropillars modulate the hybridization between the local resonances in adjacent hollow cylinders along the z direction. By tuning their radii and height, the first few photonic energy bands can be moved in frequency. We found that those energy bands mainly consist of E_z polarization (i.e., TM-like). This is consistent with the observation that Mie resonances in long hollow cylinders for TE polarization have much higher frequencies than that for the TM polarization (see Supplementary Information). Thus the degeneracy of the p - and d - bands come from space group symmetry rather than the spin degree of freedom of photon. The topological property of those photonic bands is the same as that of the energy bands of spinless bosonic systems.

We calculate the photonic energy bands for the hexagonal photonic crystal and display the results in Figs. 2a and 2b. The p and d bands cross at certain point between the Γ and A points. This crossing is accidental but unavoidable since the p and d bands are reversely ordered at the Γ and A points. The accidental degeneracy takes place at two points, $(0, 0, \pm K_z)$, which are the only p - d degeneracy points in the whole Brillouin zone (The four bands are labeled by color in Fig. 2a). For the permittivity $\varepsilon = 12$ (i.e., silicon) and the outer and inner radii, $R_{out} = 0.5a$ and $R_{in} = 0.4a$, our calculation [using MIT photonic bands (MPB)] gives $K_z = 0.34 \frac{2\pi}{h}$. The frequency of the Dirac points is $0.6 \frac{2\pi c}{a}$ (i.e., above the light line). The two points are identified as the topological Z_2 Dirac points below.

The contour surface of the electric field E_z of the p and d bands at Γ point is presented

in Figs. 2d and 2e. Their spatial symmetries can be clearly identified as p and d orbitals, which correspond to the two doubly degenerate representations of the C_{6v} symmetry group. Indeed the p bands are doubly degenerate along the Γ - A line (the same for the d bands). This degeneracy is lifted in k -space away from the Γ - A line due to pseudo-spin-orbit coupling. Also the p and d bands can mix with each other[25]. The labeling of the bands with p and d orbitals is valid only along the Γ - A line (see Fig. 2b).

The Z_2 topological number can be calculated for each k_z . Since the photonic crystal has inversion symmetry, the Z_2 topological number can be calculated by counting the parity inversion at the time-reversal invariant momenta[34]. In hexagonal photonic crystals there are only four such points for fixed k_z : three of them are equivalent to the M' point $(\frac{\pi}{a}, 0, k_z)$, the other is the Γ' point $(0, 0, k_z)$. The parity inversion only takes place at the Γ' point. If the p - and d -bands reverse order at the Γ' point (i.e., the p -bands are above the d -bands), then the Z_2 topological number is 1, otherwise it is zero. We plot the Z_2 topological number as a function of k_z in Fig. 2c. The Dirac points (its conical dispersion is shown in Fig. 2f) are then identified as the kink of the Z_2 topological number along k_z . The two Dirac points are the monopoles of the $SU(2)$ Berry-flux.

Near each $\Gamma' = (0, 0, k_z)$ point the doubly degenerate p bands can be organized as $p_+ = p_x + ip_y$ and $p_- = p_x - ip_y$ bands, which we define as the “spin-up” and “spin-down” states of the p bands. Similarly, we define the $d_+ = d_{x^2-y^2} + id_{xy}$ and $d_- = d_{x^2-y^2} - id_{xy}$ as the “spin-up” and “spin-down” states of the d bands. The coupling between those bands near the Γ' points can be obtained via a $\vec{k} \cdot \vec{P}$ theory derived from the Maxwell equations[24] (see Supplementary Information). Explicitly, the eigenvalue problem for the Maxwell equations in the photonic crystal can be written as $\nabla \times \frac{1}{\varepsilon(\vec{r})} \nabla \times \vec{\Psi}_{n,\vec{k}}(\vec{r}) = \frac{\omega_{n,\vec{k}}^2}{c^2} \vec{\Psi}_{n,\vec{k}}(\vec{r})$ (c is the speed of light in vacuum). The Hamiltonian is defined as $\mathcal{H} = \nabla \times \frac{1}{\varepsilon(\vec{r})} \nabla \times$ and the photon wavefunction for the n^{th} band with wavevector $\vec{k}$ is defined as its magnetic field, $\vec{\Psi}_{n,\vec{k}}(\vec{r}) = \vec{h}_{n,\vec{k}}(\vec{r})e^{i\vec{k} \cdot \vec{r}}$. The wavefunction (magnetic field) is normalized as $\int_{u.c.} d\vec{r} |\vec{h}_{n,\vec{k}}(\vec{r})|^2 = 1$ ($u.c.$ stands for integral in a unit cell). From the symmetry properties (C_{6v} and inversion) we found that the dominant coupling between the p - and d -bands is within the same pseudo-spin polarization (e.g., p_+ couples with d_+) (see Supplementary Information)[24, 25]. In the

basis of $(d_+, p_+, d_-, p_-)^T$ the $\vec{k} \cdot \vec{P}$ photonic Hamiltonian near the Γ' points is written as

$$\mathcal{H} = \frac{2\omega_0}{c^2} \begin{pmatrix} \frac{\omega_d^2(\vec{k})}{2\omega_0} & v_{\parallel} k_{\parallel} e^{-i\theta_k} & 0 & 0 \\ v_{\parallel}^* k_{\parallel} e^{i\theta_k} & \frac{\omega_p^2(\vec{k})}{2\omega_0} & 0 & 0 \\ 0 & 0 & \frac{\omega_d^2(\vec{k})}{2\omega_0} & v_{\parallel}^* k_{\parallel} e^{i\theta_k} \\ 0 & 0 & v_{\parallel} k_{\parallel} e^{-i\theta_k} & \frac{\omega_p^2(\vec{k})}{2\omega_0} \end{pmatrix}, \quad (1)$$

where ω_0 is the frequency of the Dirac point. To the lowest nontrivial orders in $\vec{k}_{\parallel}$ and k_z , $\omega_d(\vec{k}) = \omega_{d0}(k_z)$, $\omega_p(\vec{k}) = \omega_{p0}(k_z)$. $\theta_k = \text{Arg}(k_x + ik_y)$ and $|v_{\parallel}|$ is the velocity in the x - y plane (it is k_z -dependent). If $\omega_{p0} > \omega_{d0}$, the above Hamiltonian is similar to the Hamiltonian of the quantum spin Hall insulator[2] and hence $Z_2 = 1$. Otherwise the topology is trivial, $Z_2 = 0$. Therefore, $\omega_{p0} = \omega_{d0}$ determines a pair of kinks of the Z_2 topology. The crossing of the p and d bands at finite k_z is permitted by the fact that both p and d bands are of even-parity along z direction.

The conical dispersion (see Fig. 2f) near the Dirac points $(0, 0, \pm K_z)$ is described by the following Hamiltonian

$$\mathcal{H}_{Dirac} = \frac{\omega_0^2}{c^2} \hat{1} + \frac{2\omega_0}{c^2} \begin{pmatrix} v_d \delta k_z & v_{\parallel} k_{\parallel} e^{-i\theta_k} & 0 & 0 \\ v_{\parallel}^* k_{\parallel} e^{i\theta_k} & v_p \delta k_z & 0 & 0 \\ 0 & 0 & v_d \delta k_z & v_{\parallel}^* k_{\parallel} e^{i\theta_k} \\ 0 & 0 & v_{\parallel} k_{\parallel} e^{-i\theta_k} & v_p \delta k_z \end{pmatrix}, \quad (2)$$

where $\delta k_z = k_z - K_z$ for $k_z > 0$ (or $\delta k_z = k_z + K_z$ if $k_z < 0$). v_d and v_p are the group velocity of the p and d bands along the z direction. For k_z around K_z , $v_d > 0$ and $v_p < 0$, while these group velocities reverse sign for k_z around $-K_z$. The eigenvalue equation for the above Hamiltonian is similar to the Dirac equation for massless electrons written in the Weyl representation. Hence the Dirac points consist of a pair of Weyl points with opposite chirality. For $k_z > 0$ the p_+ and d_+ (p_- and d_-) bands form a Weyl point with chirality $+1$ (-1).

We then introduce an inversion symmetry breaking mechanism by twisting the micropillars and extend their height to $0.4a$ (see Fig. 3a-b). The resulting photonic bands along the Γ - A line (Fig. 3c) break the degeneracy between the p_+ and p_- bands as well as that between the d_+ and d_- . Hence the Dirac points are split into two pairs of Weyl points. We

find that, in consistent with the $\vec{k} \cdot \vec{P}$ theory, the crossing between the p_+ and d_+ (p_- and d_-) bands form a Weyl point with conical dispersion (Fig. 3d-e). In contrast, the crossing between p_+ and d_- (p_- and d_+) bands is quadratic. Hence there are four Weyl points along the Γ - A line (Fig. 3f).

The spatial distributions of the electromagnetic energy density for the $d_{\pm}$ and $p_{\pm}$ states are depicted in Fig. 4 (see Supplementary for more information). It is noticed that the electromagnetic energy is mostly distributed around the dielectric material (rather than in air), particularly, mainly in the hollow cylinder region rather than the micropillar region (the middle blue region in Fig. 4). The spatial patterns of the Poynting vector are also plotted in the x - y plane in Fig. 4. The Poynting vector exhibits spatial distributions similar to vortices (more precisely, Skyrmion-type patterns). The Poynting vector winds along the z axis. The winding direction is the same as the pseudo-spin direction, revealing the finite orbital angular momentum of the four states.

To show the robustness of the Dirac points, we study the dependence of the Dirac points on the inner and outer radii of the hollow cylinders [see Fig. 5a]. We find that the Dirac points emerge in a large parameter region mainly with large outer radius R_{out} . The position of the Dirac points, characterized by K_z , can be tuned by R_{in} and R_{out} . The arrows in Fig. 5a indicate the tendency that the Dirac points are moved toward the Γ point where they are created or annihilated. The robustness of the Dirac points are guaranteed by the global feature of the p - d band inversion along the Γ - A line, which is difficult to be removed by local perturbations. The surface states comprises of k_z dependent helical edge states (see Supplementary Information), connecting the two Dirac points. The parity-time symmetry and the Z_2 topology ensure that the surface states always form a closed isofrequency contour (as depicted in Fig. 5b) due to parity-time symmetry[33]. This is quite different from the Dirac points with trivial topology ($Z_2 = 0$), such as in electronic Dirac semimetals found in Ref. [35]. The trivial Dirac points are not robust and can be removed by local perturbations. Moreover, there is no surface states induced by the trivial Dirac points (see Fig. 5c).

The Weyl points emerging in inversion symmetry broken photonic crystals are also found to be stable. In Fig. 5d we give the phase diagram for the Weyl points in the R_{out} - R_{in} parameter space which is similar to that of the Dirac points. Their robustness is inherited from the Z_2 Dirac points. Fig. 5e illustrates the evolution of the nodal points and surface states upon breaking inversion symmetry. Note that the surface states connect the Weyl

points with positive and negative k_z . This is very different from the Weyl points induced by breaking inversion symmetry in systems with trivial ($Z_2 = 0$) Dirac points (Fig. 5f): The surface states connect the two Weyl points within the half-plane with $k_z > 0$ (or $k_z < 0$). The unique connectivity is a special feature of the Z_2 Dirac and Weyl points[33].

From Fig. 2b the Dirac points are above the light-line and hence accessible in transmission experiments[11]. Differing from the Dirac points, the Weyl points can only be excited by light sources with orbit angular momentum $\hbar$ or $2\hbar$ for the p_+ - d_+ Weyl points. If the orbit angular momentum reverses, they can only excite the p_- and d_- Weyl points. Thus the four Weyl point can be exploited for optical devices with efficient frequency, angle and orbit-angular-momentum filtering.

As the monopoles of the photonic $SU(2)$ gauge field, the Dirac points can be exploited to manipulate light flow based on the photonic spin Hall effect. For example, for $k_z \simeq K_z$ photons with positive orbit angular momentum ranging from $\hbar$ to $2\hbar$ (hybridization of the p_+ and d_+ orbits) develop a transvers velocity $-\Delta\vec{k} \times \vec{\Omega}(\vec{k})$ when the photonic wavevector changes $\Delta\vec{k}$ due to, e.g., scattering with an interface or an obstacle[36, 37]. The Berry-curvature is $\vec{\Omega}(\vec{k}) = \frac{\vec{q}}{q^3}$ with $\vec{q} = (k_x, k_y, k_z - K_z)$. The Berry curvature changes sign when the orbit angular momentum is reversed. In addition, the surface states can be excited at the photonic-crystal-air interface of which the velocity for positive orbit angular momentum is *opposite* to that of the negative orbit angular momentum. The abundant Berry phases and angular-momentum filtering effects enrich the manipulation of photon with finite orbit angular momentum.

This work demonstrates the atomic approach for design topological nodal points and $\vec{k}$ -space Berry phases in three-dimensional photonic crystals via Mie resonances. In addition we found a new type of Dirac points protected by the Z_2 topology and the parity-time symmetry which are monopoles of the $SU(2)$ synthetic gauge field. Our work provides new guidelines for designing topological photonic crystals for future photonic applications.

Note added.—At the final stage of this work, we noticed recent studies on Z_2 topological Dirac points in electronic energy bands[38, 39].

-
- [1] M. Z. Hasan and C. L. Kane, “Topological Insulators”, Rev. Mod. Phys. **82**, 3045 (2010).
 - [2] X.-L. Qi and S.-C. Zhang, “Topological insulators and superconductors”, Rev. Mod. Phys.

- 83**, 1057 (2011).
- [3] S. Murakami, “Phase transition between the quantum spin Hall and insulator phases in 3D: emergence of a topological gapless phase”, *New. J. Phys.* **9**, 356 (2007).
 - [4] X. Wan, A. M. Turner, A. Vishwanath, and S. Y. Savrasov, “Topological Semimetal and Fermi-Arc Surface States in the Electronic Structure of Pyrochlore Iridates”, *Phys. Rev. B* **83**, 205101 (2011).
 - [5] N. Goldman, I. Satija, P. Nikolic, A. Bermudez, M. A. Martin-Delgado, M. Lewenstein, and I. B. Spielman, “Realistic Time-Reversal Invariant Topological Insulators with Neutral Atoms”, *Phys. Rev. Lett.* **105**, 255302 (2010).
 - [6] J.-H. Jiang, “Tunable topological Weyl semimetal from simple cubic lattices with staggered fluxes”, *Phys. Rev. A* **85**, 033640 (2012).
 - [7] M. Hafezi, E. Demler, M. Lukin, and J. Taylor, “Robust optical delay lines via topological protection”, *Nat. Phys.* **7**, 907-912 (2011).
 - [8] L. Lu, L. Fu, J. D. Joannopoulos, and M. Soljačić, “Weyl points and line nodes in gapless gyroid photonic crystals”, *Nat. Photon.* **7**, 294 (2013).
 - [9] M. Hafezi, S. Mittal, J. Fan, A. Migdall, and J. Taylor, “Imaging topological edge states in silicon photonics”, *Nat. Photon.* **7**, 1001 (2013).
 - [10] L. Lu, J. D. Joannopoulos, and M. Soljačić, “Topological Photonics”, *Nat. Photon.* **8**, 821 (2014).
 - [11] L. Lu, Z. Wang, D. Ye, L. Ran, L. Fu, J. D. Joannopoulos, and M. Soljačić, “Experimental observation of Weyl points”, *Science* **349**, 622 (2015).
 - [12] W.-J. Chen, M. Xiao, and C. T. Chan, “Experimental observation of robust surface states on photonic crystals possessing single and double Weyl points”, *arXiv:1512.04681*
 - [13] K. Y. Bliokh, D. Smirnova, and F. Nori, “Quantum spin Hall effect of light”, *Science* **348**, 1448 (2015).
 - [14] Z. Yang, F. Gao, X. Shi, X. Lin, Z. Gao, Y. Chong, and B. Zhang, “Topological Acoustics”, *Phys. Rev. Lett.* **114**, 114301 (2015).
 - [15] R. Süsstrunk and S. D. Huber, “Observation of phononic helical edge states in a mechanical topological insulator”, *Science* **349**, 47 (2015).
 - [16] M. Xiao, W.-J. Chen, W.-Y. He, Z. Q. Zhang, and C. T. Chan, “Synthetic gauge fields and Weyl point in Time-Reversal Invariant Acoustic Systems”, *arXiv:1503.06295*

- [17] X. Q. Huang, Y. Lai, Z. H. Hang, H. H. Zheng, and C. T. Chan, “Dirac cones induced by accidental degeneracy in photonic crystals and zero-refractive-index materials”, *Nat. Mater.* **10**, 682 (2011).
- [18] A. Fang, Z. Q. Zhang, S. G. Louie, and C. T. Chan, “Klein tunneling and supercollimation of pseudospin-1 electromagnetic waves”, *Phys. Rev. B* **93**, 035422 (2016).
- [19] S.-L. Chua, L. Lu, J. Bravo-Abad, J. D. Joannopoulos, and M. Soljačić, “Larger-area single-mode photonic crystal surface-emitting lasers enabled by an accidental Dirac point”, *Opt. Lett.* **39**, 2072-2075 (2014).
- [20] L. Wang, S.-K. Jian, and Hong Yao, “Topological photonic crystal with equifrequency Weyl points”, arXiv:1511.09282
- [21] A. B. Khanikaev, S. H. Mousavi, W.-K. Tse, M. Kargarian, A. H. MacDonald, and G. Shvets, “Photonic topological insulators”, *Nat. Mater.* **12**, 233 (2013).
- [22] Z. Wang, Y. Chong, J. D. Joannopoulos, and M. Soljačić, “Observation of unidirectional backscattering-immune topological electromagnetic states”, *Nature (London)* **461**, 772 (2009).
- [23] M. C. Rechtsman, J. M. Zeuner, Y. Plotnik, Y. Lumer, D. Podolsky, F. Dreisow, S. Nolte, M. Segev, and A. Szameit, “Photonic Floquet topological insulators”, *Nature (London)* **496**, 196200 (2013).
- [24] K. Sakoda, “Universality of mode symmetries in creating photonic Dirac cones”, *J. Opt. Soc. Am. B* **29**, 2770 (2012).
- [25] L.-H. Wu and X. Hu, “Scheme for Achieving a Topological Photonic Crystal by Using Dielectric Material”, *Phys. Rev. Lett.* **114**, 223901 (2015)
- [26] L. Lu, C. Fang, L. Fu, S. G. Johnson, J. D. Joannopoulos, and M. Soljačić, “Symmetry-protected topological photonic crystal in three dimensions”, arXiv:1507.00337, *Nature Physics* in press.
- [27] S. John, “Strong localization of photons in certain disordered dielectric superlattices”, *Phys. Rev. Lett.* **58**, 2486 (1987).
- [28] E. Yablonovitch, “Inhibited Spontaneous Emission in Solid-State Physics and Electronics”, *Phys. Rev. Lett.* **59**, 2059 (1987).
- [29] J. D. Joannopoulos, S. G. Johnson, J. N. Winn, and R. D. Meade, *Photonic crystals: molding the flow of light* (Princeton university press, 2011).
- [30] E. Lidorikis, M. M. Sigalas, E. N. Economou, and C. M. Soukoulis, “Tight-binding

- parametrization for photonic band gap materials”, *Phys. Rev. Lett.* **81**, 1405 (1998).
- [31] L. Shi, X. Jiang, and C. Li, “Effects induced by Mie resonance in two-dimensional photonic crystals”, *J. Phys. Condens. Matter* **19**, 176214 (2007).
 - [32] L. Xu, H. X. Wang, Y. D. Xu, H. Y. Chen, and J.-H. Jiang, “Accidental degeneracy and topological phase transitions in two-dimensional core-shell dielectric photonic crystals”, *arXiv:1601.03168*
 - [33] T. Morimoto and A. Furusaki, “Weyl and Dirac semimetals with Z_2 topological charge”, *Phys. Rev. B* **89**, 235127 (2014).
 - [34] L. Fu and C. L. Kane, “Topological insulators with inversion symmetry”, *Phys. Rev. B* **76**, 045302 (2007).
 - [35] Z. K. Liu et al., “Discovery of a Three-Dimensional Topological Dirac Semimetal, Na_3Bi ”, *Science* **343**, 864 (2014).
 - [36] M. Onoda, S. Murakami, and N. Nagaosa, “Hall Effect of Light”, *Phys. Rev. Lett.* **93**, 083901 (2004)
 - [37] M. Onoda, S. Murakami, and N. Nagaosa, “Geometrical Aspects in Optical Wavepacket Dynamics”, *Phys. Rev. E* **74**, 066610 (2006).
 - [38] C. Fang, L. Lu, J. Liu, and L. Fu, “Topological semimetals with Riemann surface states”, *arXiv:1512.01552*
 - [39] B. J. Wieder, Y. Kim, A. M. Rappe, and C. L. Kane, “Double Dirac Semimetals in Three Dimensions”, *arXiv:1512.00074*

Acknowledgements

We thank supports from the National Science Foundation of China for Excellent Young Scientists (grant no. 61322504). J.H.J acknowledges supports from the faculty start-up funding of Soochow University. He also thanks Sajeev John, Suichi Murakami and Xiao Hu for helpful discussions.

Author Contributions

H.X.W performed calculations and analysis. L.X and H.Y.C contributed with advice and discussions. J.H.J conceived the idea, guided the research, and wrote the manuscript.

Materials & Correspondence

Correspondence and requests for materials should be addressed to H.Y.C (email: chy@suda.edu.cn) and J.H.J (email: joejhjiang@sina.com).

Additional information

Competing financial interests: The authors declare no competing financial interests.




FIG. 1: a, Structure in real-space unit cell of the hexagonal photonic crystals. The blue hollow cylinders and yellow micropillars are made of the same material with isotropic permittivity. b (top-down view) and c (bird's-eye view), The hexagonal photonic crystal. $\vec{a}_1$ and $\vec{a}_2$ are the two lattice vectors in the x - y plane. $a \equiv |\vec{a}_1| = |\vec{a}_2|$ is the lattice constant in the x - y plane. The height of each unit cell is $h = 0.6a$. The diameter of each micropillar is $0.1a$ and its height is $0.2a$. The outer and inner radii of the hollow cylinder is R_{out} and R_{in} . Its height is $0.4a$ d, Brillouin zone with a pair of Dirac points along the Γ - A line.




FIG. 2: a, Photonic energy band structures for the hollow-cylinder hexagonal photonic crystals with inversion symmetry (the blue curve indicate the light-line). b, The band structure along the Γ -A line. The p -bands (red) cross the d -bands (green) at $(0, 0, \pm K_z)$ with $K_z = 0.34 \frac{2\pi c}{h}$. The gray curves represent the f bands. c, The Z_2 topological number as a function of k_z . d, Isosurface plot of the E_z field of a d -orbit at the Γ point. The unit cell is depicted by the yellow dashed lines. e, Isosurface of the E_z field of a p -orbit. f, The dispersion close to the Dirac point. Parameters: $R_{out}/a = 0.5$ and $R_{in}/a = 0.4$, and $\varepsilon = 12$.




FIG. 3: Chiral hexagonal photonic crystals. a (lateral) and b (top-down) view of the structure in real-space unit cell. c, Photonic energy bands along the Γ -A line. Two Weyl points in the $k_z > 0$ region are found at $(0, 0, K_{z1})$ and $(0, 0, K_{z2})$ with $K_{z1} = 0.2 \frac{2\pi c}{h}$ and $K_{z2} = 0.26 \frac{2\pi c}{h}$. d and e, Photonic spectrum close to the two Weyl points with positive k_z . f, Depicting the four Weyl points in the whole Brillouin zone. Blue spheres denote the Weyl points with chirality -1, while the red spheres denote the Weyl points with chirality +1.




FIG. 4: Spatial field patterns of the $p_{\pm}$ and $d_{\pm}$ states at the point $\vec{k} = (0, 0, 0.1 \frac{2\pi}{h})$ in a block volume slightly larger than the unit cell of the photonic crystal. The three boundary planes (i.e., the x - y , y - z , and x - z planes) in each figure depict the energy density $\varepsilon|\vec{E}|^2(\vec{r})$ on that plane. For example, the energy density distribution at the x - y plane has a ring pattern for $p_{\pm}$ states (the ring coincides with the cross-section of the hollow cylinder). The red color represents high energy density, while the blue denote low energy density. The Poynting vector (the time-averaged value $\text{Re}[\vec{E}^* \times \vec{H}]/2$) distribution is also plotted in the x - y plane which form vortex-like patterns. The rotation directions of the Poynting vector patterns are in accordance with the pseudo-spin direction (i.e., along the z axis).




FIG. 5: a, Phase diagram of the hollow-cylinder hexagonal photonic crystal with inversion symmetry. b and c, Depicting the Dirac points and topology induced surface states in the k_z - k_x plane for systems with topological ($Z_2 = 1$) and trivial ($Z_2 = 0$) Dirac points. d, Phase diagram of the hollow-cylinder hexagonal photonic crystal with twisted micropillars (broken inversion symmetry). e and f, Illustrating the Weyl points and the topological surface states (“photon arcs”) for systems derived from the topological ($Z_2 = 1$) and trivial ($Z_2 = 0$) 3D Dirac points by introducing inversion symmetry broken perturbations. “DP” and “WP” in Figs. 5a and 5d stand for Dirac point and Weyl point, respectively.